

Scotland's Rural College

## A review of visual soil evaluation techniques for soil structure

Emmett-Booth, JP; Forristal, PD; Fenton, O; Ball, BC; Holden, NM

*Published in:*  
Soil Use and Management

*DOI:*  
[10.1111/sum.12300](https://doi.org/10.1111/sum.12300)

First published: 04/10/2016

*Document Version*  
Peer reviewed version

[Link to publication](#)

*Citation for pulished version (APA):*  
Emmett-Booth, JP., Forristal, PD., Fenton, O., Ball, BC., & Holden, NM. (2016). A review of visual soil evaluation techniques for soil structure. *Soil Use and Management*, 32(4), 623 - 634. <https://doi.org/10.1111/sum.12300>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# A Review of Visual Soil Evaluation Techniques for Soil Structure

J.P. EMMET-BOOTH<sup>1\*</sup>, P.D. FORRISTAL<sup>2</sup>, O. FENTON<sup>3</sup>, B.C. BALL<sup>4</sup> & N.M. HOLDEN<sup>1</sup>

<sup>1</sup> School of Biosystems Engineering, University College Dublin, Belfield, Dublin 4, Ireland, <sup>2</sup> Crop Science Department, Teagasc Oak Park, Co. Carlow, Ireland, <sup>3</sup>

Teagasc Environment Research Centre, Johnstown Castle, Co. Wexford, Ireland,

<sup>4</sup>SRUC Crop and Soil Systems Research Group, West Mains Road, Edinburgh, EH9 3JG, U.K.

**\*Corresponding Author:** J.P. Emmet-Booth (Email: jeremy.emmet-booth@ucdconnect.ie)

**Running Title:** Rev. of soil structure VSE Techniques

## Summary

Soil structure forms a key component of soil quality and its assessment by semi-quantitative Visual Soil Evaluation (VSE) techniques can help scientists, advisors and farmers make decisions regarding sampling and soil management. VSE techniques require inexpensive equipment and generate immediate results that correlate well with quantitative measurements of physical and biochemical properties, highlighting their potential utility. We reviewed published VSE techniques and found that soils of certain textures present problems and a lack of research into the influence of soil moisture content on VSE criteria. Generally, profile methods evaluate process interactions at the point scale, exploring both intrinsic aspects and anthropic impacts. Spade methods focus on anthropogenic characteristics, providing rapid synopses of soil structure over wider areas. Despite a focus on structural form, some methods include criteria related to stability and resiliency. Further work is needed to improve existing methods regarding texture influences, on-farm sampling procedures and more holistic assessments of soil structure.

**Keywords:** visual soil examination, soil quality, soil structure

## Introduction

Soil structure is a key component of soil quality (Askari *et al.*, 2015; Mueller *et al.*, 2013), influencing and influenced by soil chemical and biological properties (daSilva *et al.*, 2014; Askari *et al.*, 2015). Current concerns over soil resource degradation (Koch *et al.*, 2013), emphasise the importance of assessing soil structure (Mueller *et al.*, 2013). Semi-quantitative procedures for structure evaluation using visual and tactile assessment are receiving increased attention (Ball *et al.*, 2013) notably regarding impacts of current agricultural practices (Batey, 2009), reduced tillage strategies (Giarola *et al.*, 2013) and agri-environmental considerations (Newell-Price *et al.*, 2012). Soil has been visually assessed for millennia (Batey, 2000) and Visual Soil Evaluation (VSE) techniques (Ball and Munkholm, 2015) offer repeatable procedures for examining structural morphology for soil quality assessment (Mueller *et al.*, 2013). Correlations between VSE techniques and quantitative soil measurements have been widely described (McKenzie, 2001; Mueller *et al.*, 2009; Pulido Moncada *et al.*, 2014a) including indicators of soil physical quality (Guimarães *et al.*, 2013; Pulido Moncada *et al.*, 2014b) and bio-chemical quality (Askari *et al.*, 2015; daSilva *et al.*, 2014). Compared to quantitative measurements, VSE techniques provide rapid easily interpreted results using inexpensive equipment, making them widely accessible (Guimarães *et al.*, 2013; Ball *et al.*, 2007; Batey, 2000).

Due to increased interest in VSE techniques and the numerous methods in use, this review will outline the in-field procedures most widely described (according to published, English language literature from 1940 onwards), discuss VSE methodology

and synthesise strengths, weaknesses and complimentary aspects between specific procedures, thus identifying improvements.

## **An outline of in-field methods**

VSE techniques can be categorised into spade (Tables 1 and 2) and profile methods (Boizard *et al.*, 2005; Mueller *et al.* 2009). The former require soil sample blocks to be examined after extraction by spade (Tables 1 and 2), evaluating structural state up to 50cm depth, over wide areas. The latter, founded on soil survey principles (McKeague *et al.*, 1986; Batey, 2000), require examination of soil profiles to  $\approx 1.5\text{m}$  in soil pits, generally excavated mechanically (Table 3), providing detailed information at point scale.

### *Spade methods*

Görbing's (1947) Spade Diagnosis, the first published technique, focuses on anthropic impacts on structure and crop growth, qualitatively assessing soil structure, rooting and moisture content. The Peerlkamp (1959) method, the first semi-quantitative single-score procedure, together with Görbing's method, formed the foundation of more recent procedures. The Peerlkamp method focuses on anthropic impacts on structure (Boizard *et al.*, 2005). A sample block extracted to 15cm is rapidly scored between 1 and 10 considering aggregate shape, size and porosity, particle cohesion and root development. Layers can be assessed separately (Peerlkamp, 1967). The Werner Method (Werner and Thämert 1989) examines soil physical state in terms of crop growth, assessing individual layers up to 50cm. Properties including aggregate size, shape, intra-aggregate fissures, aggregate face width and bio-pores each receive a score

(Mueller *et al.*, 2009). All three methods require the manual break-up of sample blocks to expose aggregates, a key process in all spade methods.

Drop tests, which involve dropping sample blocks from a defined height onto a hard surface to expose aggregates, are also used. The Diez method (see Mueller *et al.*, 2009), first described in the late 1990's, incorporates a drop test from  $\approx 1$  m. Exposed aggregates are then assessed by hand. The method can assess anthropic impacts to 40 cm (Diez *et al.*, 2012). Soil surface condition, topsoil and subsoil structure along with rooting, redox morphology, organic matter decomposition, macro-porosity and the transition layer between topsoil and subsoil are assessed with reference to a manual, with the option of generating a single summarising score (R. Brandhuber, personal communication).

Beste (1999) developed an extended version of Görbing's Spade Diagnosis, which assesses structure for rooting and soil biota requiring the manual aggregate exposure. It incorporates a scoring system while also assessing aggregate water stability and includes quantitative measures. Layers are assessed with an emphasis on surface crust formation, silting and presence of worm casts (0-1 cm depth), aggregate shape and quantity of granular or angular aggregates (0-15 cm and 15-30 cm depth) along with aggregate shape and inter-aggregate porosity (30-40 cm depth).

Munkholm's (2000) Spade Diagnosis was founded on Preuschen's and Sobelius' Spade Diagnosis, both modified versions of Görbing's (1947) method. Layers within a sample block taken to 30 cm are examined. Boundaries, texture, aggregate type, size and grade, rupture resistance, porosity, rooting, soil fauna and organic matter

decomposition, are described in detail to support soil management. A drop test is included to determine the degree of aggregation, but most of the assessment is by manual manipulation (Table 2).

The Visual Soil Assessment (VSA) method (Shepherd 2000; 2009) captures intrinsic quality factors and anthropic impacts on soil structure. Plant and soil indicators are included and treated separately. For the main soil structure indicator, a drop test on an 8,000cm<sup>3</sup> sample block, generally extracted from the topsoil. Exposed aggregates are arranged by size on a flat surface for visual estimation of aggregate size distribution by comparison with reference photographs included in a field manual. Additional visual procedures are included for potential carbon sequestration and nutrient loss through leaching, run-off and gaseous emissions (Shepherd 2010a). For soil quality the “VS score” is the sum of individual weighted indicator scores for soil texture, structure, colour and smell, mottling, macro-porosity, the presence of earthworms, potential rooting depth, surface ponding, surface crusting and cover, and erosion.

The Soil Quality Scoring method (Ball and Douglas, 2003) is based on Munkholm’s (2000) diagnostic principles along with Beste’s (1999) scoring criteria, assesses soil physical fertility to 30cm through manual exposure of aggregates. Focusing on anthropic impacts, surface condition, soil structure (by layer) and crop rooting are assessed with reference to explanatory notes. The structure score is summed and weighted by layer depth.

The FAL Method (Hasinger *et al.*, 2004) includes a drop test on individual layers from the upper 45 cm (Boizard *et al.*, 2005). With an emphasis on anthropic impacts on

structure, aggregate size distribution is determined manually with smaller aggregates sieved from 20mm to 0.2mm. With reference to images and a coding key, aggregates are classified and the mean weight diameter and mean weight score are determined for each soil layer (Boizard *et al.*, 2005).

In 2007, three spade methods were published. The Visual Soil Structure Quality Assessment (VSSQA) described by Ball *et al.* (2007), was based on the Peerlkamp method and subsequently renamed Visual Evaluation of Soil Structure (VESS) following refinement (Guimarães *et al.*, 2011). VESS examines anthropic impacts on structure and is accessible to non-experts. From the top 25cm depth, the size, shape and visible porosity of aggregates are evaluated using an illustrated scoring key applied to individual layers. The weighted average gives an overall “*Sq*” score, similar to the system devised by Ball and Douglas (2003).

The ‘Thinksoils’ manual, developed by the United Kingdom’s Environment Agency (2007), includes instructions on conducting in-field assessment, emphasising erosion and runoff risk. Soil surface, topsoil and subsoil structure, macro-porosity, aggregate type and packing density along with plant and root growth are qualitatively examined. (Environment Agency, 2010). The M-SQR method (Mueller *et al.*, 2007) explores intrinsic soil quality and anthropic impacts to assess long-term soil quality for cropping or grazing. It is not exclusively an in-field procedure (therefore omitted from Tables 1 and 2), as regional climatic and soil survey data are incorporated (Mueller *et al.*, 2013). An overall “*SQ* score” of between 1 and 100 is generated from structural evaluation along with assessment of inherent soil properties that limit productivity (e.g. stoniness) and identifying “hazard” factors that limit soil quality (e.g. salinization).



176    *Profile methods*

177    Le Profil Cultural, originally developed by Hénin *et al.* (1960) and described by  
178    Manichon (1987), evaluates anthropic impacts (Roger-Estrade *et al.*, 2004; Peigné *et*  
179    *al.*, 2013). A soil pit the width of a seed drill ( $\approx 3\text{m} \times 1.5\text{m}$ ) is excavated perpendicular  
180    to tillage. Areas of different structure are identified in relation to horizontal layers  
181    formed by successive cultivations and lateral variation from wheels. Structure is  
182    described first by size and distribution of clods and from intra-clod porosity of clods  
183     $>2\text{mm}$ . Le Profil Cultural not only includes structural unit morphology, but also the  
184    spatial variation of overall structure. Peigné *et al.* (2013) described further steps for  
185    assessing a compact transition layer between topsoil and subsoil and biotic activity.  
186

187    The Whole Profile Method developed by Batey (2000) offers a holistic procedure for  
188    describing intrinsic soil quality and anthropic impacts on structure. The size and shape  
189    of aggregates, presence of pans, structural stability, clay mineralogy and evidence of  
190    compaction are evaluated (Batey, 2000). If required, Batey (2000) suggests the use of  
191    scoring systems described by Peerlkamp (1967) or McKenzie (1998). Principles  
192    developed by Batey form the basis of SOILpak, focused on structural characteristics  
193    associated with compaction and cotton growth (McKenzie, 1998). A soil pit, 1.5m  
194    deep and 4m long, perpendicular to tillage is recommended, from which five,  $343\text{cm}^3$   
195    samples are extracted. Scores are assigned by firmness with clod size, shape, rupture  
196    resistance, aggregation within clods and intra-clod porosity examined. If structural  
197    scores are poor, overriding factors including interconnecting porosity, smeared layers  
198    or textural changes are visually assessed. Visual assessment of aggregate stability in  
199    water is scored from a dispersion test (Field *et al.*, 1997). The SOILpak procedure has

been extended for a range of cropping systems and soils (Anderson *et al.*, 1998; McMullen, 2000).

SubVESS is an adapted version of VESS for subsoil assessment (Ball *et al.*, 2015) with an emphasis on identifying anthropic impacts on transition layers or compacted pans. Using a soil pit to 1.4m, soil layers are identified and assessed separately for mottling, strength, porosity, rooting and aggregate characteristics. “*Ssq* scores” are assigned for each layer using the SubVESS Flowchart, which provides a descriptive key and reference images. An overall *Ssq* score for the profile is expressed as a sequence of *Ssq* scores for individual layers from which any transition layer can be identified.

## **VSE Methodology**

### *Evaluation criteria*

All methods examine anthropogenic impacts on structure with some, mainly profile methods (Boizard *et al.*, 2005) additionally exploring intrinsic aspects. Aggregation (type, size and shape) and porosity form diagnostic criteria in almost all methods. Classification of the former generally assumes increased incidences of large (>5cm - >10cm), angular aggregates with higher rupture resistance, indicates poor structural quality (McKenzie, 1998; Guimarães *et al.*, 2011). Where desirable, differentiating anthropic impacts from intrinsic influences may be problematic.

Mueller *et al.* (2009) found that methods based on aggregation generated similar results, with strong correlations with measures of soil physical quality including bulk

density ( $\rho_b$ ) (Newell-Price *et al.*, 2013; Guimarães *et al.*, 2013; Mueller *et al.*, 2009), penetration resistance (Newell-Price *et al.*, 2013; Guimarães *et al.*, 2013) and air capacity (Mueller *et al.*, 2009). Additionally, VESS, largely dependent on aggregation diagnosis (Cui *et al.*, 2014), was related to soil respiration and enzyme activity (Cui *et al.*, 2015) along with chemical properties including, total carbon, soil organic carbon and total nitrogen (Askari *et al.* 2015).

Regarding aggregate determination, drop tests offer standardised, reproducible procedures of exposure. However, grass roots enmeshing aggregates (Pulido Moncada *et al.*, 2014a) and soils with high clay contents (Sonneveld *et al.*, 2014) were found to influence drop test results. Guimarães *et al.* (2011) found the manual exposure of aggregates generated the same overall results as drop-tests, despite being suggested as subjective (Ball *et al.*, 2007). Unless preformed on individual layers as the FAL method (Boizard *et al.*, 2005), drop tests do not allow the examination of aggregation within layers - a possible limitation (Giarola *et al.*, 2010; Guimarães *et al.*, 2011; Newell-Price *et al.*, 2013; Guimarães *et al.*, 2013). The delimitation of layers not only indicates potential soil functioning, but contextualises aggregation indicating anthropic influences. The evaluation of *in-situ* spatial arrangement, as employed by Le Profil Cultural, thoroughly indicates mechanisms or morphology of aggregation (Roger-Estrade *et al.*, 2004).

Both visual inter- (Shepherd 2000, Werner and Thämert 1989) and intra- (Guimarães *et al.*, 2011) aggregate porosity are examined. Exploring profile faces with a knife reveals macro-pores (Ball *et al.*, 2015), which can be highlighted with diluted paint (McKenzie, 1998). The quantification of earthworm burrows is also employed

(Munkholm, 2000; Peigné *et al.*, 2013). Mueller *et al.* (2009) found the VSA inter-aggregate porosity classification, assessed by examining an exposed face of a spade slice sample, correlated with dry  $\rho_b$ . VESS  $Sq$  scores, for which assessment of intra-aggregate porosity on exposed aggregate faces is required, corresponded with porosity determined by CT imagery (Munkholm *et al.* 2013; Garbout *et al.* 2013). The classification of clods described in Le Profil Cultural, based on intra-aggregate porosity (Peigné *et al.*, 2013), was justified with oedometer or consolidometer tests and significant differences between void ratios were reported (Roger-Estrade *et al.*, 2004). Le Profil Cultural modified for tropical soils (Neves *et al.* 2003) was found to relate to microbial biomass carbon (daSilva *et al.*, 2014).

Other criteria used include colour, redox morphology, smell and biological properties. However, techniques that include numerous, different criteria may generate the same overall result (Newel-Price *et al.*, 2013) suggesting that on certain soils, indicators additional to those centred on aggregation and porosity may be redundant. Mueller *et al.* (2009) suggested that where variation between structural states or evidence of compaction is not pronounced, procedures incorporating more diverse criteria (Shepherd *et al.*, 2000; Werner and Thämer, 1989) are desirable to achieve usable resolution. Relationships between diverse criteria and quantitative measurements were found. Pulido Moncada *et al.* (2014a) found the SQSP rooting criteria (Ball and Douglas 2003) correlated with  $\rho_b$ , soil organic carbon (SOC) and saturated hydraulic conductivity ( $K_{sat}$ ), along with the VSA soil colour criteria (Shepherd, 2009) with SOC and  $K_{sat}$ . However, the site-specific nature of such relationships is emphasised (Mueller *et al.*, 2009). Indeed VSE indicates overall structural state (Munkholm, 2000; Newel-

Price *et al.*, 2013). Universal correlation between particular quantitative measurements and VSE criteria is not necessarily expected or desirable.

In addition to structural form, surface sealing (Shepherd, 2009; Ball and Douglas 2003) or dispersion tests (Beste, 1999; McKenzie, 1998) can indicate stability, while organic matter contents, soil texture, cracking, rooting and earthworm populations (Shepherd, 2009; McKenzie, 1998) indicate resiliency, thus holistically assessing structure (Kay, 1990).

#### *Spatial, textural and moisture variation*

Profile methods are efficient at distinguishing localised variation (Roger-Estrade *et al.*, 2004; McKenzie, 2001). Spade methods being quick, though less comprehensive (Boizard *et al.*, 2013), generate accuracy through replication over wide areas. At a field scale, sampling strategies vary (Cui *et al.*, 2014; Munkholm, 2000) and further attention to on-farm procedures (Askari *et al.*, 2013) regarding survey objectives is required. Recommended minimum numbers of samples range from four for VSA (Shepherd, 2010b) to ten for VESS (Ball *et al.*, 2007) with the avoidance of damaged areas, depending on objectives (Batey, 2000). Profile method soil pit excavation is perpendicular to tillage and sufficiently long to capture damaged areas and micro-variation (McKenzie, 1998; Peigné *et al.*, 2013). Additionally, pits can be located in two contrasting areas, capturing extremes of spatial variation within a field (Ball *et al.*, 2015; McKenzie, 1998). Sampling strategies at farm scale (Sonneveld *et al.*, 2014) have received limited attention.

Texture can influence diagnostic criteria, reducing precision. Batey and McKenzie (2006) mentioned differences in cracking and rupture resistance associated with cohesive, sandy and peaty soils. Texture can be dealt with by modifying the procedure or within classification systems. A modified VSA dropt test requires sandy and loam soil samples to be dropped from 0.5m instead of 1m (Shepherd 2009). Peerlkamp (1959) described different classification systems, with the poorest class featuring dense, smooth faced aggregates on clay and loam soils, and single-grain structure on sandy soils. However, consideration must be given to agricultural management capacity, as single-grain soils when irrigated, may be highly productive. Similar classification differences were outlined by Diez *et al.* (2012), McKenzie (1998) and Ball and Douglas (2003). The latter emphasised macro-porosity and soil colour assessment in fine-textured soils, as opposed to solely aggregation.

Relationships between moisture content and VSA, SOILpak (Murphy *et al.*, 2013) and VESS (Cui *et al.* 2014) have been described. Techniques recommend deployment on moist soils (Ball *et al.*, 2007; Boizard *et al.*, 2005; Ball and Douglas, 2003, Batey, 2000), with scientific studies conducted at near Field Capacity (Abdollahi *et al.*, 2015; Pulido Moncada *et al.*, 2014b). Clearly criteria such as rupture resistance will be affected by moisture content, Munkholm (2000) and McKenzie (1998) include different diagnostic descriptions for wet and dry soils. Some older methods described by de Boodt *et al.* (1967) also include procedures for dealing with moisture content. Research on the impact of moisture content on VSE criteria and deployment is limited.

323 *VSE output*

324 Batey (2000) emphasised the importance of describing structure rather than measuring  
325 it. Qualitative outputs, generally associated with profile methods (Peigné *et al.*, 2013;  
326 Batey, 2000) - reflecting their soil survey origin, provide detailed site-specific  
327 descriptive information, potentially lost when applying numeric scores. Qualitative  
328 descriptions may not be universally comparable (Batey, 2000) though this may not be  
329 desirable. Le Profil Cultural explores point specific morphology and causes (Roger-  
330 Estrade *et al.*, 2004) not necessarily applicable elsewhere. When summarising  
331 structural state, numeric scoring systems are regarded as important (Ball *et al.*, 2015;  
332 Ball *et al.*, 2013) as they quantify structural condition, are universally comparable and  
333 allow statistical analysis (Newell-Price *et al.*, 2013; Munkholm, *et al.*, 2013). Mueller  
334 *et al.* (2009) differentiated between techniques involving the assessment of properties  
335 either, concurrently or separately. The latter (Shepherd, 2000) might enhance  
336 reliability and objectivity (Mueller *et al.*, 2009) though may not to produce a  
337 summarising score (Ball and Douglas, 2003; Munkholm, 2000; Beste, 1999; Werner  
338 and Thämert 1989).

339

340 The Peerlkamp method, a concurrent type system, generated the same overall  
341 diagnosis as a complex multi-component system (Newell-Price *et al.*, 2013). However,  
342 its ten-point scoring system is criticised as being too broad, with a five-point index  
343 identified as optimal (Ball *et al.*, 2007). This can consist of three exclusive and two  
344 intermediate classifications (Beste, 1999), or five exclusive classifications, with non-  
345 integer intermediates possible (Guimarães *et al.*, 2011). In the case of VESS, the use of  
346 integer values can limit sensitivity and interpretation (Askari *et al.*, 2013), but deci-  
347 metric scores, derived by calculation from integer values requires expert diagnosis.

Additionally, integer values can be grouped into a simple “traffic-light” colour scheme (Ball *et al.*, 2007; McKenzie, 2013), clearly indicating structural state and potential remediation requirements.

## **Strengths, Weaknesses and Complimentary Aspects**

In this section, only the most widely utilised methods are discussed.

### *Strengths and weaknesses*

VSA includes a range of intrinsic characteristics of soil quality and of structural resiliency. The VSA drop test offers a clearly defined procedure for aggregate exposure, useful for non-experts, the later modifications of which account for texture variation (Shepherd 2009), originally found to be problematic (Newell-Price *et al.*, 2013; Giarola *et al.*, 2010). However, VSA does not delimit layers. VESS considers layers and focuses on anthropic impacts, relying on the manual exposure of aggregates. This requires some experience but still is suitable for non-expert use (Ball *et al.*, 2007). Despite being reported as not dependent on texture (Cui *et al.*, 2014; Guimarães *et al.*, 2013; Giarola *et al.*, 2013; Guimarães *et al.*, 2011), VESS was problematic with fine textured soils (Askari *et al.*, 2013), an issue that Ball *et al.* (2007) originally identified (Askari *et al.*, 2015). However, Pulido Moncada *et al.* (2014a) demonstrated that VESS generated similar results to VSA while taking less time.

SOILpak examines intrinsic soil quality along with structural stability and resiliency - notably vertical porosity highlighted with paint (McKenzie, 1998). Although possibly problematic on sandier soils (Boizard *et al.*, 2005), SOILpak not only includes different scoring procedures for different textures, but also descriptions of criteria at



different moisture contents (McKenzie, 1998) while being suitable for non-expert use. In contrast, Le Profil Cultural, only applicable to arable soils, requires expertise (Roger-Estrade *et al.*, 2004) and is time consuming (Boizard *et al.*, 2005). However, it provides a comprehensive evaluation of structural morphology, notably impacts of tillage. Later descriptions (Peigné *et al.*, 2013) include criteria such as texture, cracking, and earthworm activity exploring vertical porosity, intrinsic quality, structural stability and resiliency. The analysis of clod morphology may indicate the latter (Boizard *et al.*, 2002). SubVESS, suitable for non-expert use, generates a relatively rapid evaluation of management below tillage depth. Issues may arise when differentiating anthropogenic from intrinsic features and when used on stony soils (Ball *et al.*, 2015).

#### *Complimentary aspects*

Profile methods examine point specific structural variation, assessing intrinsic quality and anthropic impacts, thus process interactions. VSA and VESS allow wider spatial evaluation and indicate structural state without thoroughly exploring mechanisms. Both approaches can be used together. SubVESS examines from 30cm depth and so should be used with VESS (Ball *et al.*, 2015). Specific technique methodology differs and can be complimentary. As Mueller *et al.* (2009) noted, where structural variation over wide areas is minimal, multi-component systems such as VSA, may be preferable over concurrent systems such as VESS. SubVESS, which places emphasis on aggregation and anthropic impacts (Ball *et al.*, 2015) worked well on a range of soil types, apart from a stony fine soil that was classified as *Ssq* 1 (good structural quality) despite being agronomically poor as indicated by Le Profil Cultural which considers intrinsic properties (Peigné *et al.*, 2013).

## Conclusion

We show wide and growing evidence of the utility of VSE techniques. An appropriate method can be selected for all situations whether research, monitoring or management. Assessment objectives, the survey area and operators' level of expertise will dictate method selection. Profile methods allow a more detailed structural assessment than spade methods, but at the cost of coverage of within-field variation due to time constraints. However, both approaches offer information not attainable using quantitative measurements. Improvements required;

- The interaction between moisture content and VSE criteria appears to have received limited attention, while variation in soil texture presents problems for some procedures. Modified procedures or classification systems according to varying textures would be of benefit, notably to VESS. Nevertheless, research shows methods are robust and valuable.
- As the utility of VSE techniques has been established, we recommended exploration of sampling strategies and analysis of spatial variation. Minimum sample replication per method should be determined.

Further research is encouraged on new procedures and on less utilised existing methods. The latter may offer useful approaches to improve more widely adopted methods and to explore wider aspects of structure such as stability and resiliency, important for an integrated and holistic assessment, notably of agricultural soils.

## Acknowledgements

This work was mainly funded by the Irish Department of Agriculture, Food and the Marine, Research Stimulus Fund grant number 13S/468.

## References

- Abdollahi, L., Hansen, E.M., Richardson, R.J. & Munkholm, L.J. 2015. Overall assessment of soil quality on humid sandy loams: Effects of location, rotation and tillage. *Soil & Tillage Research*, **145**, 29-36.
- Anderson, A., McKenzie, D. & Friend, J. 1998. *SOILpak for dryland farmers on the red soils of Central Western NSW*, NSW Agriculture, Orange. Available at: [http://www.dpi.nsw.gov.au/\\_\\_data/assets/pdf\\_file/0007/167380/soilpak-dcw-prelims.pdf](http://www.dpi.nsw.gov.au/__data/assets/pdf_file/0007/167380/soilpak-dcw-prelims.pdf) ; accessed 11/11/2015.
- Askari, M.S., Cui, J. & Holden, N.M. 2013. The visual evaluation of soil structure under arable management. *Soil & Tillage Research*, **134**, 1-10.
- Askari, M.S., Cui, J., O'Rourke, S.M. & Holden, N.M. 2015. Evaluating soil structural quality using VIS-NIR spectrum. *Soil & Tillage Research*, **146**, Part A, 108-117.
- Ball, B.C., Batey, T. & Munkholm, L.J. 2007. Field assessment of soil structural quality - a development of the Peerlkamp test. *Soil Use and Management*, **23**, 329-337.

448

449 Ball, B.C. & Douglas, J.T. 2003. A simple procedure for assessing soil structural,  
450 rooting and surface conditions. *Soil Use and Management*, **19**, 50-56.

451

452 Ball, B.C., Batey, T., Munkholm L. J., Guimarães, R.M.L., Boizard, H., McKenzie, D.  
453 C., Peigné, J., Tormena C. A. & Hargreaves P. 2015. The numeric visual  
454 evaluation of subsoil structure (SubVESS) under agricultural production. *Soil &*  
455 *Tillage Research*, **148**, 85-96.

456

457 Ball, B.C. & Munkholm, L.J. (eds). 2015. *Visual soil evaluation, realizing potential*  
458 *crop production with minimum environmental impact*. CAB International, UK.

459

460 Ball, B.C., Munkholm, L.J. & Batey, T. 2013. Applications of visual soil evaluations.  
461 *Soil & Tillage Research*, **127**, 1-2.

462

463 Batey, T. 2000. Soil profile description and evaluation. In: *Soil and environmental*  
464 *analysis, physical methods*, 2<sup>nd</sup> Ed (eds Smith K. & Mullins C) Marcel Dekker,  
465 New York, pp. 595-628.

466

467 Batey, T. 2009. Soil compaction and soil management – a review. *Soil Use and*  
468 *Management*, **25**, 335-345.

469

470 Batey, T. & McKenzie, D.C. 2006. Soil compaction: identification directly in the field.  
471 *Soil Use and Management*. **22**, 123-131.

472

- Beste, A. 1999. *An applicable field method for the evaluation of some ecologically significant soil-function-parameters in science and agricultural consulting practice*, International Soil Conservation Organisation Conference 1999. West Lafayette, Indiana USA, Institute for Soil Conservation and Sustainable Agriculture. Available et: <http://www.gesunde-erde.net/pdf-dateien/ISCO99.pdf> ; accessed 11/11/2015.
- Boizard, H., Batey, T., McKenzie, D., Rihard, G., Roger-Estrade, J., Ball, B. C., Bradley, I., Cattle, S., Hasinger, G., Munkholm, L., Murphy, B. W., Nievergelt, J., Peigné, J. & Shepherd, G. 2005. *Field meeting "visual soil structure assessment" held at the INRA Research Station, Estées Mons, France, 25-27 May 2005*. Available at; [http://iworx5.webextra.net/~istroorg/download/WG%20Visual%20Soil%20Structure%20Assessment\\_Field%20meeting.pdf](http://iworx5.webextra.net/~istroorg/download/WG%20Visual%20Soil%20Structure%20Assessment_Field%20meeting.pdf) ; accessed 11/11/2015.
- Boizard, H., Richard, G., Roger-Estrade, J., Dürr C. & Boiffin J. 2002. Cumulative effects of cropping systems on the structure of the tilled layer in northern France, *Soil & Tillage Research*, **64**, 149-164
- Boizard, H., Yoon, S.W., Leonard, J., Lheureux, S., Cousin, I., Roger-Estrade, J. & Richard, G. 2013. Using a morphological approach to evaluate the effects of traffic and weather conditions on the structure of a loamy soil in reduced tillage. *Soil & Tillage Research*, **127**, 34-44.

497 Cui, J., Askari, M.S. & Holden, H.M. 2014. Visual evaluation of soil structure under  
 498 grassland management. *Soil Use and Management*, **30**, 129-138.  
 499

500 Cui, J. & Holden, N.M. 2015. The relationship between soil microbial activity and  
 501 microbial biomass soil structure and grassland management. *Soil & Tillage*  
 502 *Research*, **146**, Part A: 32-38.  
 503

504 da Silva, A.P., Babujia, L.C., Franchini, J.C., Ralisch, R., Hungria, M. & Guimarães,  
 505 M. de F. 2014. Soil structure and its influence on microbial biomass in different  
 506 soil and crop management systems. *Soil & Tillage Research*, **142**, 42-53.  
 507

508 de Boodt M., de Leenheer L., Frese H., Lou A. & Peerlkamp P. (eds.). 1967. *West-*  
 509 *European methods for soil structure determination*. Vol **II**. The State Faculty of  
 510 Agricultural Sciences, Ghent, Belgium.  
 511

512 Diez, T., Weighlt, H. & Brandhuber, R. 2012. *Bodenstruktur erkennen und beurteilen,*  
 513 *Anleitung zur bodenuntersuchung mit dem spaten*. LfL Bayerische Landesanstalt  
 514 für Landwirtschaft, Freising-Weihenstephan. Available at:  
 515 <http://www.lfl.bayern.de/publikationen/informationen/040146/> ; accessed  
 516 11/11/2015.  
 517

518 Environment Agency. 2007. *Thinksoils manual*. Environment Agency, Bristol.  
 519 Available at:  
 520 [http://adlib.everysite.co.uk/adlib/defra/content.aspx?id=2RRVTHNXTS.892WV](http://adlib.everysite.co.uk/adlib/defra/content.aspx?id=2RRVTHNXTS.892WV1TVH19TI)  
 521 [1TVH19TI](http://adlib.everysite.co.uk/adlib/defra/content.aspx?id=2RRVTHNXTS.892WV1TVH19TI) ; accessed 11/11/2015.

522

523 Environment Agency. 2010. *Thinksoils, Examining soil structure, A practical guide to*  
524 *digging a hole*. Environment Agency, Bristol.

525

526 Field, D.J., McKenzie, D.C. & Koppi, A.J. 1997. Development of an improved  
527 Vertisol stability test for SOILpak. *Australian Journal of Soil Research*, **35**, 843-  
528 852.

529

530 Garbout, A., Munkholm, L.J. & Hansen, S.B. 2013. Tillage effect on topsoil structural  
531 quality assessed using X-ray CT soil cores and visual soil evaluation. *Soil &*  
532 *Tillage Research*, **128**, 104-109.

533

534 Giarola, N.F.B., da Silva, A.P., Tormena, C.A., Ball, B. & Rosa, J.A. 2010. Visual soil  
535 structure quality assessment on Oxisols under no-tillage system. *Scientia*  
536 *Agricola*, **64**, 479-482.

537

538 Giarola, N.F.B., da Silva, A.P., Tormena, C.A., Guimarães, R.M.L. & Ball, B.C. 2013.  
539 On the visual evaluation of soil structure: The Brazilian experience in Oxisols  
540 under no-tillage. *Soil & Tillage Research*, **127**, 60-64.

541

542 Görbing, J. 1947. *Die grundlagen der gare im praktischen ackerbau, Band II*.  
543 Landbuch- Verlang G.M.B.H., Hannover.

544

545 Guimarães, R.M.L., Ball, B.C., Tormena, C.A., Giarola, N.F.B. & da Silva, A.P. 2013.  
546 Relating visual evaluation of soil structure to other physical properties in soils of  
547 contrasting texture and management. *Soil & Tillage Research*, **127**, 92-99.  
548

549 Guimarães, R.M.L., Ball, B.C. & Tormena, C.A. 2011. Improvements in the visual  
550 evaluation of soil structure. *Soil Use and Management*, **27**, 395-403.  
551

552 Hasinger, G., Nievergelt, J., Petrasek, M. & Weisskopf, P. 2004. *Observer et évaluer*  
553 *la structure du sol*. Les catiers de le FAL 50, Agroscope.  
554

555 Hénin *et al.* 1960. WAITING FOR INTER-LIBRARY LOAN  
556

557 Kay B.D. 1990. Rates of change of soil structure under different cropping systems. In:  
558 *Advances in soil science* (ed Stewart B.A.), Springer, New York, 12, pp. 1-52.  
559

560 Koch, A., McBratney, A., Adams, M., Field, D., Hill, R., Crawford, J., Minasny, B.,  
561 Lal, R., Abbott, L., O'Donnell, A., Angers, D., Baldock, J., Barbier, E., Binkly,  
562 D., Parton, W., Wall, D.H., Bird, M., Bouma, J., Chenu, C., Flora, C.B.,  
563 Grunwold, S., Hempel, J., Jastrow, J., Lehmann, J., Lorenz, K., Morgan, C.,  
564 Rice, C., Whitehead, D., Yang, I. & Zimmermann, M. 2013. Soil security:  
565 solving the global soil crisis. *Global Policy*, **4**, 434-441.  
566

567 Manichon, H. 1987. Observation morphologique de l'état structural et mise en  
568 évidence d'effets de compactage des horizons travaillés. In: *Soil compaction and*  
569 *regeneration, proceedings of the workshop on soil compaction: consequences*



570        *and structural regeneration processes, Avignon 17-18 September 1985* (eds.  
571        Monnier, G. & Goss, M.J.), AA Balkema, Rotterdam, Boston, pp. 39-52  
572

573        McKeague, J.A., Wang, C. & Coen, G.M. 1986. *Describing and interpreting the*  
574        *macrostructure of mineral soils – a preliminary report*. Land Resource Research  
575        Institute Ottawa, Ontario, LRRRI Contribution No. 84-50.  
576

577        McKenzie, D. 1998. *SOILpak for cotton growers*, 3<sup>rd</sup> Ed. NSW Agriculture, Orange.  
578        <http://www.dpi.nsw.gov.au/agriculture/resources/soils/guides/soilpak/cotton> ;  
579        accessed 11/11/2015.  
580

581        McKenzie, D.C. 2001. Rapid assessment of soil compaction damage II. Relationships  
582        between the SOILpak score, strength and aeration measurements, clod shrinkage  
583        parameters and image analysis data on a vertisol. *Australian Journal of Soil*  
584        *Research*, **39**, 127-141.  
585

586        McKenzie, D.C. 2013. Visual soil examination techniques as part of a soil appraisal  
587        framework for farm evaluations in Australia. *Soil & Tillage Research*, **127**, 26-  
588        33.  
589

590        McMullen, B. 2000. *SOILpak for vegetable growers*. NSW Agriculture, Orange.  
591        Available at:  
592        [http://www.dpi.nsw.gov.au/\\_\\_data/assets/pdf\\_file/0020/127208/contents.pdf](http://www.dpi.nsw.gov.au/__data/assets/pdf_file/0020/127208/contents.pdf) ;  
593        accessed 11/11/2015.  
594

595 Mueller, L., Kay, B.D., Hu, C., Li, Y., Schindler, U., Behrendt, A., Shepherd, T.G. &  
 596 Ball, B. C., 2009. Visual assessment of soil structure: Evaluation of  
 597 methodologies on sites in Canada, China and Germany, Part I: Comparing visual  
 598 methods and linking them with soil physical data and grain yield of cereals. *Soil*  
 599 *& Tillage Research*, **103**, 178-187.

600

601 Mueller, L., Schindler, U., Behrendt, A., Eulenstein, F. & Dannowski, R. 2007. *The*  
 602 *Muencheberg soil quality rating (SQR), Field manual for detecting and*  
 603 *assessing properties and limitations of soils for cropping and grazing*, Leibniz-  
 604 Centre for Agriculture Landscape Research (ZALF) e. V. Muencheberg  
 605 Germany. Available at:  
 606 [http://www.zalf.de/de/forschung/institute/lwh/mitarbeiter/lmueller/Documents/field\\_mueller.pdf](http://www.zalf.de/de/forschung/institute/lwh/mitarbeiter/lmueller/Documents/field_mueller.pdf) ; accessed 11/11/2015.

607

608

609 Mueller, L., Shepherd, G., Schindler, V., Ball, B.C., Munkholm, L.J., Hennings, V.,  
 610 Smolentseve, E., Rukhovic, O., Lukin, S. & Hu, C. 2013. Evaluation of soil  
 611 structure in the framework of an overall soil quality rating. *Soil & Tillage*  
 612 *Research*, **127**, 74-84.

613

614 Munkholm, L.J. 2000. *The spade analysis – a modification of the qualitative spade*  
 615 *diagnosis for scientific use*. DIAS Report April 2000 No. 28, Plant Production.  
 616 Danish Institute of Agricultural Sciences, Denmark.

617

618 Munkholm, L.J., Heck, R.J. & Deen, B. 2013. Long-term rotation and tillage effects on  
 619 soil structure and crop yield. *Soil & Tillage Research*, **127**, 85-91

- Murphy, B.W., Crawford, M.H., Duncan, D.A., McKenzie, D.C. & Koen, T.B. 2013. The use of visual soil assessment schemes to evaluate surface structure in a soil monitoring program. *Soil & Tillage Research*, **127**, 3-12.
- Neves, C.S.V.J., Feller, C., Guimarães, M.F., Medina, C.C., Tavares Filho, J. & Fortier, M. 2003. Soil bulk density and porosity of homogeneous morphological units identified by the Cropping Profile Method in clayey Oxisols in Brazil. *Soil & Tillage Research*, **71**, 109-119
- Newell-Price, J.P., Chambers, B. & Whittingham, M. 2012. BD5001: Characterisation of soil structural degradation under grassland and development measures to ameliorate its impact on biodiversity and other soil functions. Grassland Soil Compaction Assessment – Stages 1 and 2, May 2012. ADAS, U.K. Available at: <http://randd.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None&ProjectID=16827&FromSearch=Y&Publisher=1&SearchText=BD5001&SortString=ProjectCode&SortOrder=Asc&Paging=10#Description;> accessed 08/03/2016.
- Newell-Price, J.P., Whittingham, M.J., Chambers, B.J. & Peel, S. 2013. Visual soil evaluation in relation to measured soil physical properties in a survey of grassland soil compaction in England and Wales. *Soil & Tillage Research*, **127**, 65-73.

645 Peerlkamp, P. 1959. *A visual method of soil structure evaluation*. Meded. v.d.  
646 Landbouwhogeschool en Opzoekingsstations van der Staat te Gent. XXIV No.  
647 24, pp. 216-221  
648

649 Peigné, J., Vian, J-F, Cannavacciuola, M., Lefeure, V., Gautronneau, Y. & Boizard, H.  
650 2013. Assessment of soil structure in the transition layer between topsoil and  
651 subsoil using the profil cultural method. *Soil & Tillage Research*, **127**, 13-25.  
652

653 Pulido Moncada, M., Gabriels, D., Lobo, D., Rey, J.C. & Cornelis, W.M. 2014a.  
654 Visual field assessment of soil structural quality in tropical soils. *Soil & Tillage*  
655 *Research*, **139**, 8-18.  
656

657 Pulido Moncada, M., Penning, L. H., Timm, L. C., Gabriels, D. & Cornelis, W. M.  
658 2014b. Visual examination and soil physical and hydraulic properties for  
659 assessing soil structural quality of soils with contrasting textures and land uses.  
660 *Soil & Tillage Research*, **140**, 20-28.  
661

662 Roger-Estrade, J., Richard, G., Caneill, J., Boizard, H., Defosse, P. & Manichon, H.  
663 2004. Morphological characterisation of soil structure in tilled fields: from  
664 diagnosis methods to modelling of structural changes over time. *Soil & Tillage*  
665 *Research*, **79**, 33-49.  
666

667 Shepherd, T.G. 2000. *Visual soil assessment. Volume 1. Field guide for cropping and*  
668 *pastoral grazing on flat to rolling country*. horizons.mw & Landcare Research,  
669 Palmerston North, New Zealand. Available at:

670 [http://www.landcareresearch.co.nz/\\_\\_data/assets/pdf\\_file/0011/28676/VSA\\_Vol](http://www.landcareresearch.co.nz/__data/assets/pdf_file/0011/28676/VSA_Vol)  
 671 [ume1\\_smaller.pdf](#) ; accessed 11/11/2015.  
 672  
 673 Shepherd, T.G. 2009. *Visual soil assessment, Field guide for pastoral grazing and*  
 674 *cropping on flat to rolling country*. 2<sup>nd</sup> Ed. Horizons Regional Council, New  
 675 Zealand.  
 676  
 677 Shepherd, T.G. 2010a. *Visual soil assessment, Pastures, Part 2, Visual indicators of*  
 678 *environmental performance under pastoral grazing*. Food and Agriculture  
 679 Organisation, Rome. Available at:  
 680 <http://www.fao.org/docrep/010/i0007e/i0007e00.htm>; accessed 11/11/2015.  
 681  
 682 Shepherd, T.G. 2010b. *Visual soil assessment, Pastures Part 1*. Food and Agricultural  
 683 Organisation, Rome. Available at:  
 684 <http://www.fao.org/docrep/010/i0007e/i0007e00.htm>; accessed 18/03/2016  
 685  
 686 Sonneveld, M.P.W., Heuvelink, G.B.M. & Moolenaar, S.W., 2014. Application of  
 687 visual soil examination and evaluation technique at site and farm level. *Soil Use*  
 688 *and Management*, **30**, 263-271.  
 689  
 690 Werner, D. & Thämert, W. 1989. *Zur diagnose des physikalischen bodenzustandes auf*  
 691 *produktionsflächen*. Arch. Acker-Pflanzenbau Bodenkd, Berlin. **33**, 729-739.  
 692

693  
694  
695

Table 1 Outline of VSE Spade Methods (Drop Test Procedures)

Method*	Origin	Objective	Land assessed	Characteristics assessed	Criteria employed	Depth assessed	Scoring system used	Intended users	Time requirement
The Diez Method (Diez <i>et al.</i> , 2012)	Germany	To assess structure in relation to soil functioning, notably plant growth and water infiltration	Emphasis on arable	Anthropic impacts on structure	Aggregate type, size, shape, inter-aggregate porosity, rooting, redox morphology, transition layer	40 <sup>b</sup> cm	Score between 1 and 5 used ( <i>1 = best, 5 = worst</i> )	Advisors and farmers <sup>b</sup>	- -
Visual Soil Assessment (VSA) (Shepherd, 2000, 2009, 2010)	New Zealand	To assess soil state, plant performance and the impact of farm management	Arable and grassland	Intrinsic soil quality and anthropic impacts on structure	Texture, aggregate size distribution, macro-porosity, redox morphology surface ponding and deformation, earthworms, smell, colour, potential rooting depth	Varying depths	VS score of between 0 and 50 ( <i>&lt;20 = poor, 20-35 = moderate, &gt;35 = good</i> )	Advisors and farmers	40 minutes
FAL Method (Hasinger <i>et al.</i> , 2004)	Switzerland	To provide an accurate evaluation of structural state at a specific point <sup>a</sup>	Arable and grassland	Anthropic impacts on structure	Aggregate type, size, distribution and mean weight diameter <sup>a</sup>	45 cm	Score between 1 and 14 used for aggregate mean score ( <i>1 = worst, 14 = best</i> ). Aggregate mean weight diameter is described in mm	Researchers and advisors <sup>a</sup>	90 minutes <sup>a</sup>

\*Sources provided are not necessarily the original description of methods, <sup>a</sup> Sourced from: Boizard *et al.* (2005), <sup>b</sup> Points are of the authors' opinions

696  
697

698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723

724  
725

**Table 2** Outline of VSE Spade Methods (Manual Aggregate Exposure Procedures)

Method*	Origin	Objective	Land assessed	Characteristics assessed	Criteria employed	Depth assessed	Scoring system used	Intended users	Time requirement
Spade Diagnosis (Görbing 1947)	Germany	To assess structure in relation to plant growth	Emphasis on arable	Anthropic impacts on structure	Aggregate size, shape, porosity and rooting	30 cm	No numeric scores used	Advisors and farmers	- -
Peerlkamp Method (Peerlkamp, 1959)	The Netherlands	To assess structure in relation to fertility, summarised by a single score	Emphasis on arable	Anthropic impacts on structure	Aggregate size shape, rupture resistance, inter- and intra-porosity, rooting, surface soil dispersion	15 cm	<i>St</i> Score between 1 and 10 ( <i>1 = worst, 10 = best</i> )	Researchers, advisors and farmers <sup>a</sup>	30 minutes for 10 assessments <sup>a</sup>
The Werner Method (Werner and Thämer 1989)	Germany	To assess soil physical condition in relation to plant growth	- -	Anthropic impacts on structure	Layers, aggregate size, width, shape, inter-aggregate porosity, bio-pores	50 cm	Scores between 1 and 4 or 1 and 5 used to describe individual properties, resulting in a five digit nominal value score for each layer	Researchers	- -
Extended Spade Diagnosis (Beste, 1999)	Germany	To assess structure with regard to rooting conditions	Emphasis on arable	Anthropic impacts on structure	Aggregate type, size, shape, along with aggregate	40 cm	Scores between 1 and 5 used for structure and between	Advisors and farmers	- -



		and habitats for soil biota			stability		0 and 2 for silting type. Three sample layers are assessed separately		
Spade Analysis (Munkholm, 2000)	Denmark	To describe and relate soil tilth to management while aiding and evaluating soil management decisions	Emphasis on arable	Intrinsic soil quality and anthropic impacts on structure	Texture, colour, layer boundaries, aggregate size, shape, grade, soil consistence, macro-porosity, pore distribution, connectivity, orientation and rooting, OM decomposition and soil fauna	30 cm	Different scoring systems used for different properties, though no summarising numeric scores used	Researchers and advisors <sup>a</sup>	1 – 3 hours <sup>a</sup>
Soil Quality Scoring Procedure (SQSP) (Ball and Douglas, 2003)	United Kingdom	To assess physical fertility in terms of structure, rooting and soil surface conditions	Arable and grassland	Anthropic impacts on structure	Soil surface, aggregate type, size, shape, rupture resistance and rooting	30 cm	Three separate scores are assigned, each between 1 and 5 ( <i>1 = worst, 5 = best</i> )	Researchers and advisors <sup>a</sup>	1 hour <sup>a</sup>
Visual Soil Structure Quality Assessment (VSSQA) –	United Kingdom	To semi-quantitatively assess soil structural quality in a	Arable and grassland	Anthropic impacts on structure	Aggregate size, shape, intra-porosity, rupture	25 cm	<i>Sq</i> Score between 1 and 5 ( <i>1 = best, 5 = worst</i> )	Advisors and farmers	15 minutes

Visual Evaluation of Soil Structure (VESS) (Guimarães <i>et al.</i> , 2011)		manner accessible to non-experts			resistance rooting, redox- morphology				
Thinksoils Manual (Environment Agency, 2007, 2010)	United Kingdom	To assess soil structure with regard to erosion and run-off potential	Arable and grassland	Anthropic impacts on structure	Fissures and porosity, aggregate size, shape, rupture resistance, redox morphology, rooting, crop growth	40 cm	No numeric scores used	Advisors and farmers	- -

\*Sources provided are not necessarily the original description of methods, <sup>a</sup> Sourced from: Boizard *et al.* (2005)

730  
731

**Table 3** Outline of VSE Profile Methods

Method*	Origin	Objective	Land assessed	Characteristics assessed	Criteria employed	Depth assessed	Scoring system used	Intended users	Time requirement
Le Profil Cultural (Peigné <i>et al.</i> , 2013)	France	To examine the impact of tillage on soil structure features	Arable	Emphasis on anthropic impacts on structure	Soil layers, structural zones, macro-pores, aggregate/clod size, intra-porosity, redox morphology, rooting	1.5 m	No numeric score used	Researchers	1 – 3 hours <sup>a</sup>
Whole Profile Assessment (Batey, 2000)	United Kingdom	To assess the anthropic impact on intrinsic soil properties in relation to crop growth <sup>a</sup>	Arable and grassland	Intrinsic soil quality and anthropic impacts on structure	Soil layers, texture, aggregate size, shape, aggregate stability, compacted zones, soil bearing capacity, soil colour, redox morphology	1.2 - 1.5 m	No numeric score used	Researchers and consultants <sup>a</sup>	20 – 40 minutes <sup>a</sup>
SOILpak (McKenzie, 1998)	Australia	To identify and assess compaction in relation to crop growth	Emphasis on arable	Intrinsic soil quality and anthropic impacts on structure	Texture, soil surface, rooting, aggregate size, shape, rupture resistance, macro-pores, aggregate stability	1.5 m	Score between 0 and 2 used for structural ( <i>0 = worst, 2 = best</i> ) and ASWAT score	Land surveyors, advisors and farmers	25 – 90 minutes <sup>a</sup>

							between 0 and 16 used for aggregate stability ( <i>0 = negligible dispersion, 16 = serious dispersion</i> )	
SubVESS Flowchart (Ball <i>et al.</i> , 2015)	United Kingdom	To assesses any anthropogenic transition layer in terms of crop growth	Emphasis on arable	Anthropic impacts on structure	Redox morphology, porosity, rooting, aggregate size, shape	1.4 m	<i>Ssq</i> scores of between 1 and 5 ( <i>1 = best, 5 = worst</i> )	Advisors 20 minutes

\*Sources provided are not necessarily the original description of methods, <sup>a</sup> Sourced from: Boizard *et al.* (2005)